

# New results on dark energy in the early and present Universe and the observational status of the inflationary scenario

Alexei A. Starobinsky

Landau Institute for Theoretical Physics RAS,  
Moscow - Chernogolovka, Russia

Theoretical seminar  
IPP RAS, Moscow, 19.06.2014

Present status of inflation

Anomalous growth of light scalar fields in the de Sitter space-time

Inflationary spectral predictions and observations

Outcome of recent CMB observations

Consequences of the primordial GW discovery

Conclusions

# Four epochs of the history of the Universe

$H \equiv \frac{\dot{a}}{a}$  where  $a(t)$  is a scale factor of an isotropic homogeneous spatially flat universe (a Friedmann-Lemaître-Robertson-Walker background):

$$ds^2 = dt^2 - a^2(t)(dx^2 + dy^2 + dz^2).$$

The history of the Universe in one line according to the present paradigm:

$$? \longrightarrow DS \Longrightarrow FLWRD \Longrightarrow FLWMD \Longrightarrow \overline{DS} \longrightarrow ?$$

$$|\dot{H}| \ll H^2 \Longrightarrow H = \frac{1}{2t} \Longrightarrow H = \frac{2}{3t} \Longrightarrow |\dot{H}| \ll H^2$$

$$p \approx -\rho \Longrightarrow p = \rho/3 \Longrightarrow p \ll \rho \Longrightarrow p \approx -\rho$$

# Main advantages of inflation

## Aesthetic elegance

Inflation – hypothesis about an almost maximally symmetric (quasi-de Sitter) stage of the evolution of our Universe in the past, before the hot Big Bang. If so, preferred initial conditions for (quantum) inhomogeneities with sufficiently short wavelengths exist – the adiabatic in-vacuum ones. In addition, these initial conditions represent an attractor for a much larger compact open set of initial conditions having a non-zero measure in the space of all initial conditions.

## Predictability, proof and/or falsification

Given equations, this gives a possibility to calculate all subsequent evolution of the Universe up to the present time and even further to the future. Thus, any concrete inflationary model can be proved or disproved by observational data.

## Naturalness of the hypothesis

Remarkable qualitative similarity between primordial and present dark energy.

# Present status of inflation

## From "proving" inflation to using it as a tool

Present status of inflation: transition from "proving" it in general and testing some of its simplest models to applying the inflationary paradigm to investigate particle physics at super-high energies and the actual history of the Universe in the remote past using real observational data on  $n_s(k) - 1$  and  $r(k)$ .

# Generation of scalar and tensor perturbations during inflation

A genuine quantum-gravitational effect: a particular case of the effect of particle-antiparticle creation by an external gravitational field. Requires quantization of a space-time metric. Similar to electron-positron creation by an electric field. From the diagrammatic point of view: an imaginary part of a one-loop correction to the propagator of a gravitational field from all quantum matter fields including the gravitational field itself, too.

The effect can be understood from the behaviour of a light scalar field in the de Sitter space-time.

# De Sitter space-time

Constant curvature space-time.

$$R_{\alpha\beta\gamma\delta} = H_0^2 (g_{\alpha\gamma} g_{\beta\delta} - g_{\alpha\delta} g_{\beta\gamma})$$

4 most popular forms of its space-time metric (only the first metric covers the whole space-time):

$$ds^2 = dt_c^2 - H_0^{-2} \cosh^2(H_0 t_c) (d\chi_c^2 + \sin^2 \chi_c d\Omega^2)$$

$$ds^2 = dt^2 - a_1^2 e^{2H_0 t} (dr^2 + r^2 d\Omega^2), \quad a_1 = \text{const}$$

$$ds^2 = dt_o^2 - H_0^{-2} \sinh^2(H_0 t_o) (d\chi_o^2 + \sinh^2 \chi_o d\Omega^2)$$

$$ds^2 = (1 - H_0^2 R^2) d\tau^2 - (1 - H_0^2 R^2)^{-1} dR^2 - R^2 d\Omega^2$$

$$d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$$

# Perturbative anomalous growth of light scalar fields in the de Sitter space-time

Background - **fixed** - de Sitter or, more interestingly, quasi-de Sitter space-time (slow roll inflation).

Occurs for  $0 \leq m^2 \ll H^2$  where  $H \equiv \frac{\dot{a}}{a}$ ,  $a(t)$  is a FRW scale factor. The simplest and textbook example:

$m = 0$ ,  $H = H_0 = \text{const}$  for  $t \geq t_0$  and the initial quantum state of the scalar field at  $t = t_0$  is the adiabatic vacuum for modes with  $k/a(t_0) \gg H_0$  and some infrared finite state otherwise.

The wave equation:

$$\phi_{;\mu}^{;\mu} = 0$$

Equation for the time dependent part of tensor perturbations on a FLRW background supported by ideal fluids or minimally coupled scalar fields has the same form.



Quantization with the adiabatic vacuum initial condition:

$$\phi = (2\pi)^{-3/2} \int \left[ \hat{a}_{\mathbf{k}} \phi_{\mathbf{k}}(\eta) e^{-i\mathbf{k}\mathbf{r}} + \hat{a}_{\mathbf{k}}^\dagger \phi_{\mathbf{k}}^* e^{i\mathbf{k}\mathbf{r}} \right] d^3k$$

$$\phi_{\mathbf{k}}(\eta) = \frac{H_0 e^{-i\mathbf{k}\eta}}{\sqrt{2k}} \left( \eta - \frac{i}{k} \right), \quad a(\eta) = \frac{1}{H_0 \eta}, \quad \eta_0 < \eta < 0, \quad k = |\mathbf{k}|$$

Then

$$\langle \phi^2 \rangle = \frac{H_0^2 N}{4\pi^2} + \text{const}$$

Here  $N = \ln \frac{a}{a(t_0)} \gg 1$  is the number of e-folds from the beginning of inflation and the constant depends on the initial quantum state (Linde, 1982; AS, 1982; Vilenkin and Ford, 1982).

Straightforward generalization to the slow-roll case  $|\dot{H}| \ll H^2$ .

For  $0 < m^2 \ll H^2$ , the Bunch-Davies equilibrium value

$$\langle \phi^2 \rangle = \frac{3H_0^4}{8\pi m^2} \gg H_0^2$$

is reached after a large number of e-folds  $N \gg \frac{H_0^2}{m^2}$ .  
Purely infrared effect - creation of real field fluctuations;  
renormalization is not important and does not affect it.

For the de Sitter inflation (gravitons only) (AS, 1979):

$$P_g(k) = \frac{16GH_0^2}{\pi}; \quad \langle h_{ik}h^{ik} \rangle = \frac{16GH_0^2 N}{\pi}.$$

The assumption of small perturbations breaks down for  $N \gtrsim 1/GH_0^2$ . Still ongoing discussion on the final outcome of this effect. My opinion - no screening of the background cosmological constant, instead - stochastic drift through an infinite number of locally de Sitter, but globally non-equivalent vacua.

Reason: the de Sitter space-time is not the generic late-time asymptote of classical solutions of GR with a cosmological constant  $\Lambda$  both without and with hydrodynamic matter. The generic late-time (expanding) asymptote is (AS, 1983):

$$ds^2 = dt^2 - \gamma_{ik} dx^i dx^k$$

$$\gamma_{ik} = e^{2H_0 t} a_{ik} + b_{ik} + e^{-H_0 t} c_{ik} + \dots$$

where  $H_0^2 = \Lambda/3$  and the matrices  $a_{ik}$ ,  $b_{ik}$ ,  $c_{ik}$  are functions of spatial coordinates.  $a_{ik}$  contains two independent physical functions (after 3 spatial rotations and 1 shift in time + spatial dilatation) and can be made unimodular, in particular.

# Generation of metric perturbations

One spatial Fourier mode  $\propto e^{i\mathbf{k}\mathbf{r}}$  is considered.

For scales of astronomical and cosmological interest, the effect of creation of metric perturbations occurs at the primordial de Sitter (inflationary) stage when  $k \sim a(t)H(t)$  where  $k \equiv |\mathbf{k}|$  (the first Hubble radius crossing).

After that, for a very long period when  $k \ll aH$  until the second Hubble radius crossing (which occurs rather recently at the radiation or matter dominated stages), there exist one mode of scalar (adiabatic, density) perturbations and two modes of tensor perturbations (primordial gravitational waves) for which metric perturbations are constant (in some gauge) and independent of (unknown) local microphysics due to the causality principle.

# Classical-to-quantum transition for the leading modes of perturbations

In the superhorizon regime in the coordinate representation:

$$ds^2 = dt^2 - a^2(t)(\delta_{lm} + h_{lm})dx^l dx^m, \quad l, m = 1, 2, 3$$

$$h_{lm} = 2\zeta(\mathbf{r})\delta_{lm} + \sum_{a=1}^2 g^{(a)}(\mathbf{r}) e_{lm}^{(a)}$$

$$e_l^{l(a)} = 0, \quad g^{(a)}_{,l} e_m^{l(a)} = 0, \quad e_{lm}^{(a)} e^{lm(a)} = 1$$

$\zeta$  describes primordial scalar perturbations,  $g$  – primordial tensor perturbations (primordial gravitational waves (GW)).

Quantum-to-classical transition: in fact, metric perturbations  $h_{lm}$  are quantum (operators in the Heisenberg representation) and remain quantum up to the present time. But, after omitting of a very small part, decaying with time, they become commuting and, thus, equivalent to classical (c-number) stochastic quantities with the Gaussian statistics (up to small terms quadratic in  $\zeta, g$ ).

Remaining quantum coherence: deterministic correlation between  $\mathbf{k}$  and  $-\mathbf{k}$  modes - shows itself in the appearance of acoustic oscillations (primordial oscillations in case of GW).

# FLRW dynamics with a scalar field

In the absence of spatial curvature and other matter:

$$H^2 = \frac{\kappa^2}{3} \left( \frac{\dot{\phi}^2}{2} + V(\phi) \right)$$

$$\dot{H} = -\frac{\kappa^2}{2} \dot{\phi}^2$$

$$\ddot{\phi} + 3H\dot{\phi} + V'(\phi) = 0$$

where  $\kappa^2 = 8\pi G$  ( $\hbar = c = 1$ ).

# Inflationary slow-roll dynamics

Slow-roll occurs if:  $|\ddot{\phi}| \ll H|\dot{\phi}|$ ,  $\dot{\phi}^2 \ll V$ , and then  $|\dot{H}| \ll H^2$ .

Necessary conditions:  $|V'| \ll \kappa V$ ,  $|V''| \ll \kappa^2 V$ . Then

$$H^2 \approx \frac{\kappa^2 V}{3}, \quad \dot{\phi} \approx -\frac{V'}{3H}, \quad N \equiv \ln \frac{a_f}{a} \approx \kappa^2 \int_{\phi_f}^{\phi} \frac{V}{V'} d\phi$$

First obtained in [A. A. Starobinsky, Sov. Astron. Lett. 4, 82 \(1978\)](#) in the  $V = \frac{m^2 \phi^2}{2}$  case and for a bouncing model.



# Spectral predictions of the one-field inflationary scenario in GR

Scalar (adiabatic) perturbations:

$$P_{\zeta}(k) = \frac{H_k^4}{4\pi^2 \dot{\phi}^2} = \frac{GH_k^4}{\pi |\dot{H}|_k} = \frac{128\pi G^3 V_k^3}{3 V_k'^2}$$

where the index  $k$  means that the quantity is taken at the moment  $t = t_k$  of the Hubble radius crossing during inflation for each spatial Fourier mode  $k = a(t_k)H(t_k)$ . Through this relation, the number of e-folds from the end of inflation back in time  $N(t)$  transforms to  $N(k) = \ln \frac{k_f}{k}$  where  $k_f = a(t_f)H(t_f)$ ,  $t_f$  denotes the end of inflation.  
The spectral slope

$$n_s(k) - 1 \equiv \frac{d \ln P_{\zeta}(k)}{d \ln k} = \frac{1}{k^2} \left( 2 \frac{V_k''}{V_k} - 3 \left( \frac{V_k'}{V_k} \right)^2 \right)$$

Generically  $n_s \neq 1$ , though  $|n_s - 1| \ll 1$  – deviation from the Harrison-Zeldovich spectrum is expected!

The special case when  $n_s \equiv 1$ :  $V(\phi) \propto \phi^{-2}$  in the slow-roll approximation.

Omitting the slow-roll assumption:

let  $x = \sqrt{4\pi G}\phi$ ,  $y = B\sqrt{4\pi G}H$ ,  $v(x) = \frac{32\pi^2 G^2 B^2}{3} V(\phi)$ .

Then (A. A. Starobinsky, JETP Lett. 82, 169 (2005)):

$$y = e^{x^2/2} \left( \int_x^\infty e^{-\tilde{x}^2/2} d\tilde{x} + C \right)$$

$$v = y^2 - \frac{1}{3} \left( \frac{dy}{dx} \right)^2$$

Tensor perturbations (A. A. Starobinsky, JETP Lett. 50, 844 (1979)):

$$P_g(k) = \frac{16GH_k^2}{\pi}; \quad n_g(k) \equiv \frac{d \ln P_g(k)}{d \ln k} = -\frac{1}{k^2} \left( \frac{V'_k}{V_k} \right)^2$$

The consistency relation:

$$r(k) \equiv \frac{P_g}{P_\zeta} = \frac{16|\dot{H}_k|}{H_k^2} = 8|n_g(k)|$$

Tensor perturbations are always **suppressed** by at least the factor  $\sim 8/N(k)$  compared to scalar ones. For the present Hubble scale,  $N(k_H) = (50 - 60)$ .

# From metric perturbations to CMB anisotropy and polarization

## I. Temperature anisotropy

For scalar perturbations, generated mainly at the last scattering surface at  $z_{LSS} \approx 1090$  (the Sachs-Wolfe, Silk and Doppler effects), but also after it (the integrated Sachs-Wolfe effect).

For GW – only the ISW works.

## II. Polarization

Produced at the last scattering surface only due to the Thomson scattering of photons on electrons, suppressed by the factor  $\Delta z_{LSS}/z_{LSS} \sim 0.1$  compared to a temperature anisotropy.

No circular polarization, only linear one.

Linear polarization on the sky (2-sphere) can be decomposed into the E-mode (scalar) and the B-mode (pseudoscalar).

1. Expand the  $Q \pm iU$  combinations of the Stokes parameters into spin-weighted spherical harmonics  $_{\pm 2}Y_{lm}$ .
2. Then

$$a_{E,lm} = -(a_{2,lm} + a_{-2,lm}) / 2, \quad a_{B,lm} = (a_{2,lm} - a_{-2,lm}) / 2$$

In the first order, the E-mode is produced both by scalar perturbations and GW, the B-mode is produced by GW only. The most important second order effect through which scalar perturbations produce B-mode: gravitational lensing of CMB fluctuations, screens the first order effect for multipoles  $l > 150$ .

# Outcome of recent CMB observations

## I. A year ago

The most important for the history of the early Universe are:

1. The primordial spectrum of scalar perturbations has been measured and its deviation from the flat spectrum  $n_s = 1$  in the first order in  $|n_s - 1| \sim N^{-1}$  has been discovered (using the multipole range  $\ell > 40$ ):

$$P_\zeta(k) = \int \frac{\Delta_\zeta^2(k)}{k} dk, \quad \Delta_\zeta^2 = (2.20^{+0.05}_{-0.06}) 10^{-9} \left(\frac{k}{k_0}\right)^{n_s-1}$$

$$k_0 = 0.05 \text{Mpc}^{-1}, \quad n_s - 1 = -0.040 \pm 0.007$$

N.B.: The value is obtained under some natural assumptions, the most critical of them is  $N_\nu = 3$ , for  $N_\nu = 4$  many things have to be reconsidered and  $n_s \approx 1$  is not excluded.

2. Neither the B-mode of CMB polarization, nor primordial GW were discovered:  $r < 0.11$  at the 95% CL.

II. Three months ago


BISEP2 collaboration: P. A. R. Ade et al., arXiv:1403.3985:  
discovery of the B-mode in the multipole range  $30 < l < 150$   
(for larger  $l$  it was discovered earlier this year with the amount  
in agreement from gravitational lensing of scalar  
perturbations) with

$$r = 0.20_{-0.05}^{+0.07}$$


The unsubtracted result – contains an unknown foreground non-thermal part.

Consequence – assuming the Einstein gravity:

$$\sqrt{G} H_{dS} = 0.99 \times 10^{-5} \left( \frac{r_{0.002}}{0.2} \right)^{1/2} 5^{0.96-n_s}$$



-200  $T(\mu\text{K})$  +200 WMAP 5-year






# Combined results from Planck and other experiments



P. A. R. Ade et al., arXiv:1303.5082

Model	Parameter	Planck+WP	Planck+WP+lensing	Planck + WP+high- $\ell$	Planck+WP+BAO
$\Lambda$ CDM + tensor	$n_s$	$0.9624 \pm 0.0075$	$0.9653 \pm 0.0069$	$0.9600 \pm 0.0071$	$0.9643 \pm 0.0059$
	$r_{0.002}$	$< 0.12$	$< 0.13$	$< 0.11$	$< 0.12$
	$-2\Delta \ln \mathcal{L}_{\text{max}}$	0	0	0	-0.31

**Table 4.** Constraints on the primordial perturbation parameters in the  $\Lambda$ CDM+ $r$  model from *Planck* combined with other data sets. The constraints are given at the pivot scale  $k_* = 0.002 \text{ Mpc}^{-1}$ .



**Fig. 1.** Marginalized joint 68% and 95% CL regions for  $n_s$  and  $r_{0.002}$  from *Planck* in combination with other data sets compared to the theoretical predictions of selected inflationary models.



# Consequences of the primordial GW discovery

If confirmed by an independent measurement:

1. Discovery of a real singularity – a state of the Universe in the past with a very high curvature (with  $H$  only 5 orders of magnitude less than the Planck mass).
2. Discovery of a new class of gravitational waves – primordial ones.
3. Decisive argument for the necessity of quantization of gravitational waves.
4. Decisive test of the inflationary paradigm as a whole.
5. Discovery of  $\sim 20\%$  deviation of the power spectrum of scalar perturbations from a scale-free one – new physics during inflation!

The most intriguing discordance between WMAP and Planck results from one side and the BISEP2 ones from the other: **no sign** of GW in the CMB temperature anisotropy spectrum.

Instead of the  $1 + \frac{6.20}{8}r \approx 1.1$  increase of the total anisotropy power spectrum over the multipole range  $2 \ll l < 50$ , a  $\sim 10\%$  depression is seen for  $20 \lesssim l \lesssim 40$  (see e.g. Fig. 39 of arXiv:1303.5076).

6.20 is the rounded value for

$$\frac{25}{9} \left( 1 + \frac{48\pi^2}{385} \right)$$

(A. A. Starobinsky, *Sov. Astron. Lett.* 11, 133 (1985)). In this expression, only the Sachs-Wolfe effect is taken into account for scalar perturbations. Adding the Doppler and Silk effects leads to the decrease of this number to  $\approx 5$  for the range  $20 \lesssim l \lesssim 40$ .

More detailed analysis in D. K. Hazra, A. Shafieloo, G. F. Smoot and A. A. Starobinsky, arXiv:1403.7786 : the power-law form of  $P_\zeta(k)$  is excluded at more than  $3\sigma$  CL.

# Broken scale models describing both WMAP-Planck and BISEP2 data



Next step: "whipped inflation" D. K. Hazra, A. Shafieloo, G. F. Smoot and A. A. Starobinsky, arXiv:1404.0360.

The model contains a new scale at which the effective inflaton potential has a feature which the inflaton crosses about 50 e-folds before the end of inflation. The existence of such a feature, in turn, requires some new physics (e.g. fast phase transition in a second field coupled to the inflaton).

$$V(\phi) = V_S(\phi) + V_R(\phi)$$

$$V_S(\phi) = \gamma\phi^p, \quad V_R(\phi) = \lambda(\phi - \phi_0)^q\Theta(\phi - \phi_0)$$

Best results for  $(p, q) = (2, 3)$ .



# Wiggles in the power spectrum

The effect of the **same order**: an upward wiggle at  $l \approx 40$  and a downward one at  $l \approx 22$ .

**Lesson**: irrespective of a future analysis of foreground contamination in the BISEP2 result, features in the anisotropy spectrum for  $20 \lesssim l \lesssim 40$  confirmed by WMAP and Planck should be taken into account and studied seriously.

A more elaborated class of model suggested by previous studies of sharp features in the inflaton potential caused, e.g. by a fast phase transition occurred in another field coupled to the inflaton during inflation:

**WWI** (Wiggly Whipped inflation)

D. K. Hazra, A. Shafieloo, G. F. Smoot and A. A. Starobinsky,  
[arXiv:1405.2012](https://arxiv.org/abs/1405.2012)



In particular, the potential with a sudden change of its first derivative:

$$V(\phi) = \gamma\phi^2 + \lambda\phi^p(\phi - \phi_0)\theta(\phi - \phi_0)$$

which generalizes the exactly soluble model considered in A. A. Starobinsky, JETP Lett. **55**, 489 (1992) produces  $-2\Delta \ln \mathcal{L} = -11.8$  compared to the best-fitted power law scalar spectrum, partly due to the better description of wiggles at both  $l \approx 40$  and  $l \approx 22$ .

A sharp feature in the potential leads to a rapid increase of the effective inflaton mass,  $m^2 = V''(\phi)$ , in the vicinity of  $\phi = \phi_0$ . While  $m \approx 2 \times 10^{13}$  GeV for  $\phi < \phi_0$ , it becomes of the order of  $10^{14}$  GeV and larger at earlier times when  $\phi \geq \phi_0$  (but still much less than the energy density scale of the inflaton potential  $\sim 3 \times 10^{16}$  GeV).

# Conclusions

- ▶ Inflation is being transformed into a normal physical theory, based on some natural assumptions confirmed by observations and used to obtain new theoretical knowledge from them.
- ▶ First **quantitative** observational evidence for small quantities of the first order in the slow-roll parameters:  $n_s(k) - 1$  and  $r(k)$ .
- ▶ The quantitative theoretical prediction of these quantities is based on gravity (space-time metric) quantization and requires very large space-time curvature in the past of our Universe with a characteristic length only five orders of magnitude larger than the Planck one.
- ▶ The BISEP2 result by itself is the confirmation of the general prediction (made in 1979) of the early Universe scenario with the de Sitter (inflationary) stage preceding the radiation dominated stage (the hot Big Bang).

- ▶ However, would the BISEP2 result be confirmed, inflation is not so simple: the scalar primordial power spectrum deviates from a scale-free one that implies the existence of some scale (i.e. new physics) during inflation.
- ▶ Though the Einstein gravity plus a minimally coupled inflaton remains sufficient for description of inflation with existing observational data, modified (in particular, scalar-tensor or  $f(R)$ ) gravity can do it as well.
- ▶ The conceptual change in utilizing CMB and other observational data from "proving inflation" to using them to determine the spectrum of particle masses in the energy range  $(10^{13} - 10^{14})$  GeV by making a "tomographic" study of inflation.